

Semileptonic Hyperon Decays and CKM Unitarity*

Nicola Cabibbo

*Department of Physics, University of Rome - La Sapienza
and INFN, Sezione di Roma 1
Piazzale A. Moro 5, 00185 Rome, Italy
nicola.cabibbo@roma1.infn.it*

Earl C. Swallow

*Department of Physics, Elmhurst College
Elmhurst, Illinois 60126 and Enrico Fermi Institute,
The University of Chicago, Chicago, Illinois
earls@elmhurst.edu*

Roland Winston

*Division of Natural Sciences,
The University of California - Merced,
Merced, California 95344
rwinston@ucmerced.edu
(Dated: February 7, 2008)*

Using a technique that is not subject to first-order $SU(3)$ symmetry breaking effects, we determine the V_{us} element of the CKM matrix from data on semileptonic hyperon decays. We obtain $V_{us} = 0.2250(27)$, where the quoted uncertainty is purely experimental. This value is of similar experimental precision to the one derived from K_{l3} , but it is higher and thus in better agreement with the unitarity requirement, $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$. An overall fit including the axial contributions, and neglecting $SU(3)$ breaking corrections, yields $F + D = 1.2670 \pm 0.0035$ and $F - D = -0.341 \pm 0.016$ with $\chi^2 = 2.96/3$ d.f.

PACS numbers: 12.15.Hh, 13.30.Ce, 14.20.Jn

The determination of the elements of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [1, 2] is one of the main ingredients for evaluating the solidity of the standard model of elementary particles. This is a vast subject which has seen important progress with the determination [3, 4] of ϵ'/ϵ and the observation [5, 6] of CP violation in B decays.

While a lot of attention has recently been justly devoted to the higher mass sector of the CKM matrix, it is the low mass sector, in particular V_{ud} and V_{us} where the highest precision can be attained. The most sensitive test of the unitarity of the CKM matrix is provided by the relation $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1 - \Delta$. Clearly the unitarity condition is $\Delta = 0$. The $|V_{ub}|^2$ contribution [7] is negligible (10^{-5}) at the current level of precision. The value $V_{ud} = 0.9740 \pm 0.0005$ is obtained from superallowed pure Fermi nuclear decays [8]. In combination with $V_{us} = 0.2196 \pm 0.0023$, derived from K_{e3} decay [9, 10], this yields $\Delta = 0.0032 \pm 0.0014$. On its face, this represents a 2.3 standard deviation departure from unitarity [8].

In this communication we reconsider the contribution that the hyperon beta decays can give to the determination of V_{us} . The conventional analysis of hyperon beta decay in terms of the parameters F, D and V_{us} is marred

by the expectation of first order $SU(3)$ breaking effects in the axial-vector contribution. The situation is only made worse if one introduces adjustable $SU(3)$ breaking parameters as this increases the number of degrees of freedom and degrades the precision. If on the contrary, as we do here, one focuses the analysis on the vector form factors, treating the rates and g_1/f_1 [11] as the basic experimental data, one has directly access to the f_1 form factor for each decay, and this in turn allows for a redundant determination of V_{us} . The consistency of the values of V_{us} determined from the different decays is a first confirmation of the overall consistency of the model. A more detailed version of this work will be published in the Annual Reviews of Nuclear and Particle Sciences [12].

In 1964 Ademollo and Gatto proved [13] that there is no first-order correction to the vector form factor, $\Delta^1 f_1(0) = 0$. This is an important result: since experiments can measure $V_{us} f_1(0)$, knowing the value of $f_1(0)$ in $\Delta S = 1$ decays is essential for determining V_{us} .

The Ademollo-Gatto Theorem suggests an analytic approach to the available data that first examines the vector form factor f_1 because it is not subject to first-order $SU(3)$ symmetry breaking effects. An elegant way to do this is to use the *measured* value of g_1/f_1 along with the predicted values of f_1 and f_2 to extract a V_{us} value from the decay rate for each decay. If the theory is correct, these should coincide within errors, and can be combined to obtain a best value of V_{us} . This consistency of the V_{us}

*To be published in Physical Review Letters.

values obtained from different decays then indicates the success of the Cabibbo model. A similar approach appears to have been taken in Ref. [14].

Four hyperon beta decays have sufficient data to perform this analysis: $\Lambda \rightarrow p e^- \bar{\nu}$, $\Sigma^- \rightarrow n e^- \bar{\nu}$, $\Xi^- \rightarrow \Lambda e^- \bar{\nu}$, $\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}$ [9]. Table I shows the results for them. In this analysis, both model-independent and model-dependent radiative corrections [11] are applied and q^2 variation of f_1 and g_1 is included. Also $SU(3)$ values of $g_2 = 0$ and f_2 are used along with the numerical rate expressions tabulated in Ref. [11]. We have not however included $SU(3)$ -breaking corrections to the f_1 form factor, which will be discussed in the next section. The stated V_{us} errors are purely experimental, coming from experimental uncertainties in the hyperon lifetimes, branching ratios, and form factor ratios.

The four values are clearly consistent ($\chi^2 = 2.26/3\text{d.f.}$) with the combined value of $V_{us} = 0.2250 \pm 0.0027$. This value is nearly as precise as that obtained from kaon decay ($V_{us} = 0.2196 \pm 0.0023$) and, as observed in previous analyses [15, 16, 26], is somewhat larger. In combination with $V_{ud} = 0.9740 \pm 0.0005$ obtained from superallowed pure Fermi nuclear decays [8], the larger V_{us} value from hyperon decays beautifully satisfies the unitarity constraint $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$.

TABLE I: Results from V_{us} analysis using measured g_1/f_1 values

Decay Process	Rate (μsec^{-1})	g_1/f_1	V_{us}
$\Lambda \rightarrow p e^- \bar{\nu}$	3.161(58)	0.718(15)	0.2224 ± 0.0034
$\Sigma^- \rightarrow n e^- \bar{\nu}$	6.88(24)	-0.340(17)	0.2282 ± 0.0049
$\Xi^- \rightarrow \Lambda e^- \bar{\nu}$	3.44(19)	0.25(5)	0.2367 ± 0.0099
$\Xi^0 \rightarrow \Sigma^+ e^- \bar{\nu}$	0.876(71)	$1.32(+.22/- .18)$	0.209 ± 0.027
Combined	—	—	0.2250 ± 0.0027

We will limit our discussion to the effects that are most relevant for the determination of V_{us} . Turning our attention first to $SU(3)$ -breaking corrections to the f_1 form factor, we find in the literature computations that use some version of the quark model, as in [17, 18], or some version of chiral perturbation theory, as in [15, 19, 20].

The quark-model computations find that the f_1 form factors for the different $\Delta S = 1$ decays are reduced by a factor, the same for all decays, given as 0.987 in [17], and 0.975 in [18], a decrease respectively of 1.3% or 2.5%. This is a very reasonable result, the decrease arising from the mismatch of the wave functions of baryons containing different numbers of the heavier s quarks.

Evaluations of f_1 in chiral perturbation theory range from small negative corrections in [19] to larger positive corrections in [15, 20]. Positive corrections in f_1 for *all* hyperon beta decays cannot be excluded, but are certainly not expected in view of an argument [21] according to which one expects a negative correction to f_1 at least in the $\Sigma^- \rightarrow n e^- \bar{\nu}$ case. This result follows from the observation that the intermediate states that con-

tribute to the positive second-order terms in the Ademollo and Gatto sum rule have, in this case, quantum numbers $S = -2$, $I = 3/2$; no resonant baryonic state is known with these quantum numbers. If we accept the hypothesis that the contribution of resonant hadronic states dominate, we can conclude that the correction to f_1 in Σ^- beta decay should be negative. We note that this argument also applies to K_{l3} decays, and that the corrections to these decays, computed with chiral perturbation theory, are, as expected, negative.

A modern revisitation of the quark-model computations will be feasible in the near future with the technologies of lattice QCD, and we would expect that a small negative correction would be obtained in *quenched* lattice QCD, an approximation that consists in neglecting components in the wave function of the baryons with extra quark-antiquark pairs. This is known to be an excellent approximation in low-energy hadron phenomenology [22].

Multiquark effects can be included in lattice QCD by forsaking the quenched approximation for a *full* simulation. Alternatively one could resort to chiral perturbation theory to capture the major part of the multiquark contributions which will be dominated by virtual π , K , η states. Early results of a similar strategy applied to the K_{e3} decays [23] indicate that in that case a 1% determination of the $f_+(0)$ form factor is within reach, and we expect that a similar precision can be obtained in the case of hyperon decays. In the present situation we consider it best not to include any $SU(3)$ breaking corrections in our evaluation, nor to include an evaluation of a theoretical error. Our expectation that the corrections to $f_+(0)$ will be small and negative can only be substantiated by further work.

We next turn our attention to the possible effect of ignoring the g_2 form factor. In the absence of second class currents [24] the form factor g_2 can be seen to vanish in the $SU(3)$ symmetry limit. The argument is very straightforward: the neutral currents $A_\alpha^3 = \bar{q}\lambda^3\gamma_\alpha\gamma_5q$ and $A_\alpha^8 = \bar{q}\lambda^8\gamma_\alpha\gamma_5q$ that belong to the same octet as the weak axial current are even under charge conjugation, so that their matrix elements cannot contain a weak – electricity term, which is C -odd. The vanishing of the weak electricity in the proton and neutron matrix elements of A_α^3 , A_α^8 implies the vanishing of the D and F coefficients for $g_2(0)$, so that, in the $SU(3)$ limit, the $g_2(0)$ form factor vanishes for any current in the octet.

In hyperon decays a nonvanishing $g_2(0)$ form factor can arise from the breaking of $SU(3)$ symmetry. Theoretical estimates [25] indicate a value for $g_2(0)/g_1(0)$ in the -0.2 to -0.5 range.

In determining the axial-vector form factor g_1 from the Dalitz Plot — or, equivalently, the electron–neutrino correlation — one is actually measuring $\tilde{g}_1$, a linear combination of g_1 and g_2 ($\tilde{g}_1 \approx g_1 - \delta g_2$ up to first order in $\delta = \Delta M/M$). This has already been noticed in past experiments and is well summarized in Gaillard and Sauvage [26], Table 8. Therefore, in deriving $V_{us}^2 f_1^2$ (hence V_{us}) from the beta decay rate, there is in fact a

small sensitivity to g_2 . To first order, the rate is proportional to $V_{us}^2[f_1^2 + 3g_1^2 - 4\delta g_1 g_2] \approx V_{us}^2[f_1^2 + 3\tilde{g}_1^2 + 2\delta \tilde{g}_1 g_2]$. In fact, this is a second order correction to the value of V_{us} , potentially of the same order of magnitude as the corrections to f_1 .

Experiments that measure correlations with polarization — in addition to the electron–neutrino correlation — are sensitive to g_2 . While the data are not yet sufficiently precise to yield good quantitative information, one may nevertheless look for trends. In polarized $\Sigma^- \rightarrow p e^- \bar{\nu}$ [27], negative values of g_2/f_1 are clearly disfavored (a positive value is preferred by 1.5σ). Since the same experiment unambiguously established that g_1/f_1 is negative one concludes that allowing for nonvanishing g_2 would increase the derived value of $V_{us}^2 f_1^2$. In polarized $\Lambda \rightarrow p e^- \bar{\nu}$ the data favor [28] negative values of g_2/f_1 (by about 2σ). In this decay, g_1/f_1 is positive so that again, allowing for the presence of nonvanishing g_2 would increase the derived value of $V_{us}^2 f_1^2$. In either case, we may conclude that making the conventional assumption of neglecting the g_2 form factor tends to *underestimate* the derived value of V_{us} . A more quantitative conclusion must await more precise experiments. We consider it to be of the highest priority to determine the g_2 form-factor (or a stringent limit on its value) in at least one of the hyperon decays, ideally in Λ semileptonic decay which at the moment seems to offer the single most precise determination of V_{us} .

The excellent agreement with the unitarity condition of our determination of V_{us} , which neglects $SU(3)$ -breaking effects, seems to indicate that such effects were overestimated in the past, probably as a consequence of the uncertainties of the early experimental results. We also find [12] that the g_1 form factor of the different decays, which

is subject to first order corrections, is well fitted by the F, D parameters [1], with $F + D = 1.2670 \pm 0.0035$ and $F - D = -0.341 \pm 0.016$ with $\chi^2 = 2.96/3$ d.f.

The value of V_{us} obtained from hyperon decays is of comparable experimental precision with that obtained from K_{l3} decays, and is in better agreement with the value of θ_C obtained from nuclear beta decay. While a discrepancy between V_{us} and V_{ud} could be seen as a portent of exciting new physics, a discrepancy between the two different determinations of V_{us} can only be taken as an indication that more work remains to be done both on the theoretical and the experimental side.

On the theoretical side, renewed efforts are needed for the determination of $SU(3)$ -breaking effects in hyperon beta decays as well as in K_{l3} decays. While it is quite possible to improve the present situation on the quark-model front, the best hopes lie in lattice QCD simulations, perhaps combined with chiral perturbation theory for the evaluation of large-distance multi-quark contributions.

We have given some indication that the trouble could arise from the K_{l3} determination of V_{us} , and we would like to encourage further experimental work in this field [29]. We are however convinced of the importance of renewed experimental work on hyperon decays, of the kind now in progress at the CERN SPS. The interest of this work goes beyond the determination of V_{us} , as it involves the intricate and elegant relationships that the model predicts.

The continuing intellectual stimulation provided by colleagues in the Fermilab KTeV Collaboration, particularly members of the hyperon working group, is gratefully acknowledged. This work was supported in part by the U.S. Department of Energy under grant DE-FG02-90ER40560 (Task B).

-
- [1] N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963)
 - [2] M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
 - [3] KTeV Collaboration, A. Alavi-Harati *et al.*, *Phys. Rev. D* **67**:012005 (2003)
 - [4] NA48 Collaboration, A. Lai *et al.*, *Eur. Phys. J. C* **22**, 231 (2001) [arXiv:hep-ex/0110019].
 - [5] BABAR Collaboration, B. Aubert *et al.*, *Phys. Rev. Lett.* **87**, 091801 (2001).
 - [6] Belle Collaboration, K. Abe *et al.*, *Phys. Rev. Lett.* **87**, 091802 (2001).
 - [7] BABAR Collaboration, B. Aubert *et al.*, *Phys. Rev. Lett.* **90**, 181801 (2003).
 - [8] I. S. Towner and J. C. Hardy, Santa Fe 1998, *Physics beyond the standard model*, 338-359; [arXiv:nucl-th/9809087].
 - [9] K. Hagiwara *et al.* (Particle Data Group), *Phys. Rev. D* **66**:010001 (2002)
 - [10] H. Leutwyler and M. Roos, *Z. Phys. C* **25**, 91 (1984)
 - [11] A. Garcia and P. Kielanowski, *The Beta Decay of Hyperons*, Lecture Notes in Physics **222**, Springer-Verlag, Berlin, 1985. With the metric and γ matrix conventions used here, g_1/f_1 is **positive** for neutron beta decay.
 - [12] N. Cabibbo, E. C. Swallow, R. Winston, *Ann. Rev. Nucl. Part. Sci.* Dec. 2003, **53**: 39-75.
 - [13] M. Ademollo and R. Gatto, *Phys. Rev. Lett.* **13**, 264 (1964).
 - [14] P. M. Gensini and G. Violini, hep-ph/9311270.
 - [15] R. Flores-Mendieta, E. Jenkins and A. V. Manohar, “SU(3) symmetry breaking in hyperon semileptonic decays,” *Phys. Rev. D* **58**, 094028 (1998) [arXiv:hep-ph/9805416].
 - [16] R. Flores-Mendieta, A. Garcia, G. Sanchez-Colon *Phys. Rev. D* **54**:6855 (1996);
 - [17] J. F. Donoghue, B. R. Holstein and S. W. Klimt, “K-M Angles And SU(3) Breaking In Hyperon Beta Decay,” *Phys. Rev. D* **35**, 934 (1987).
 - [18] F. Schlumpf, “Beta decay of hyperons in a relativistic quark model,” *Phys. Rev. D* **51**, 2262 (1995) [arXiv:hep-ph/9409272].
 - [19] A. Krause, “Baryon Matrix Elements Of The Vector Current In Chiral Perturbation Theory,” *Helv. Phys. Acta* **63**, 3 (1990).
 - [20] J. Anderson and M. A. Luty, “Chiral corrections to hy-

- peron vector form-factors,” Phys. Rev. D **47**, 4975 (1993) [arXiv:hep-ph/9301219].
- [21] H. R. Quinn and J. D. Bjorken, “Renormalization Of Weak Form-Factors, And The Cabibbo Angle,” Phys. Rev. **171**, 1660 (1968).
 - [22] J. Sexton, D. Weingarten, *Phys. Rev. D* **55**:4025 (1997)
 - [23] We are grateful to Guido Martinelli for a discussion on this point, see D. Bećirević *et al.*, [arXiv:hep-ph/0403217].
 - [24] S. Weinberg, Phys. Rev. **112**, 1375 (1958).
 - [25] Barry R. Holstein, “Hyperon 99”, Proceedings of the Hyperon Physics Symposium, pgs. 4-9, Fermilab, September 27-29 1999, D.A. Jensen, E. Monnier, Editors.
 - [26] J. M. Gaillard and G. Sauvage, Ann. Rev. Nucl. Part. Sci. **34**, 351 (1984).
 - [27] S. Y. Hsueh, *et al.*, Phys. Rev. D **38**, 2056 (1988).
 - [28] R. Oehme, R. Winston, A. Garcia, Phys. Rev. D **3**, 1618 (1971).
 - [29] In fact, a recently reported result from the Brookhaven E865 Collaboration indicates a higher K_{e3} decay rate than that previously used to determine V_{us} . See A. Sher, *et al.*, hep-ex/0305042 (June 24, 2003).